Published by Institute of Physics Publishing for SISSA/ISAS
Received:
Revised:
Accepted:

arXiv:hep-th/0608052v2 6 Oct 2006

JHEP00(2002)000

Odd Nambu bracket on Grassmann algebra

Dmitrij V. Soroka and Vyacheslav A. Soroka

Kharkov Institute of Physics and Technology
1 Akademicheskaya St., 61108 Kharkov, Ukraine
E-mail: `dsoroka@kipt.kharkov.ua`, `vsoroka@kipt.kharkov.ua`

Abstract: The Grassmann-odd Nambu bracket on the Grassmann algebra is proposed.

Keywords: Odd Nambu Bracket, BRST Symmetry.

Dedicated to Valentina Erofeyevna Korol'.

© SISSA/ISAS 2008

Contents

1. Introduction 1

2. Odd Nambu bracket 2

3. Conclusion 3

1. Introduction

In the paper [1] the linear odd bracket corresponding to the $SO(3)$ group and realized solely on the Grassmann variables θ_α has been introduced (see also [2, 3, 4, 5])

$$\{A,B\}_1 = A \overleftarrow{\partial}_{\theta_\alpha} \epsilon_{\alpha\beta\gamma}\theta_\gamma \overrightarrow{\partial}_{\theta_\beta} B, \quad (\alpha,\beta,\gamma = 1,2,3), \tag{1.1}$$

where $\epsilon_{\alpha\beta\gamma}$ is the completely antisymmetric Levi-Civita tensor, $\overleftarrow{\partial}$ and $\overrightarrow{\partial}$ are the right and left derivatives and $\partial_{\theta_\alpha} \equiv \frac{\partial}{\partial\theta_\alpha}$. It was constructed for this bracket three Grassmann-odd nilpotent Δ-like differential operators of the first, second and third orders with respect to the Grassmann derivatives

$$\Delta_{+1} = \epsilon_{\alpha\beta\gamma}\theta_\alpha\theta_\beta\partial_{\theta_\gamma}, \tag{1.2}$$

$$\Delta_{-1} = \frac{1}{2}\epsilon_{\alpha\beta\gamma}\theta_\alpha\partial_{\theta_\beta}\partial_{\theta_\gamma}, \tag{1.3}$$

$$\Delta_{-3} = \frac{1}{3!}\epsilon_{\alpha\beta\gamma}\partial_{\theta_\alpha}\partial_{\theta_\beta}\partial_{\theta_\gamma}. \tag{1.4}$$

The operator Δ_{+1} is proportional to the second term in a BRST charge

$$Q = \theta_\alpha G_\alpha - \frac{1}{2}\epsilon_{\alpha\beta\gamma}\theta_\alpha\theta_\beta\partial_{\theta_\gamma},$$

where θ_α and ∂_{θ_α} represent the operators for the ghosts and ghost momenta respectively and G_α are generators of the $SO(3)$ group. The operator Δ_{-1} related to the divergence of the vector field $\{\theta_\alpha, A\}_1$

$$\Delta_{-1}A = -\frac{1}{2}\partial_{\theta_\alpha}\{\theta_\alpha, A\}_1 \tag{1.5}$$

is proportional to the true Δ-operator for the odd bracket (1.1). The operator Δ_{-1} (1.5) determines the linear odd bracket (1.1) as a deviation of the Leibniz rule under the usual multiplication

$$\Delta_{-1}(A\cdot B) = (\Delta_{-1}A)\cdot B + (-1)^{p(A)}A\cdot\Delta_{-1}B + (-1)^{p(A)}\{A,B\}_1\ .$$

and simultaneously satisfies the Leibniz rule with respect to the linear odd bracket composition

$$\Delta_{-1}(\{A, B\}_1) = \{\Delta_{-1}A, B\}_1 + (-1)^{p(A)+1}\{A, \Delta_{-1}B\}_1,$$

where $p(A)$ is a Grassmann parity of the quantity A.

In the present paper we show that the operator Δ_{-3} (1.4) is related with the Grassmann-odd Nambu bracket [6] on the Grassmann algebra.

2. Odd Nambu bracket

By applying the operator Δ_{-3} (1.4) to the usual product of two quantities A and B, we obtain

$$\Delta_{-3}(A \cdot B) = (\Delta_{-3}A) \cdot B + (-1)^{p(A)}A \cdot \Delta_{-3}B + (-1)^{p(A)}\Delta(A, B), \tag{2.1}$$

where the quantity $\Delta(A, B)$ is

$$\Delta(A, B) = \frac{1}{2}\epsilon_{\alpha\beta\gamma}\left[\left(\partial_{\theta_\alpha}\partial_{\theta_\beta}A\right)\partial_{\theta_\gamma}B + (-1)^{p(A)}\left(\partial_{\theta_\alpha}A\right)\partial_{\theta_\beta}\partial_{\theta_\gamma}B\right]. \tag{2.2}$$

By acting with the operator Δ_{-3} on the usual product of three quantities A, B and C, we come to the following relation:

$$\begin{aligned}\Delta_{-3}(A \cdot B \cdot C) = {} & (\Delta_{-3}A) \cdot B \cdot C + (-1)^{p(A)}A \cdot (\Delta_{-3}B) \cdot C + (-1)^{p(A)+p(B)}A \cdot B \cdot \Delta_{-3}C \\ & + (-1)^{p(A)}\Delta(A, B)C + (-1)^{p(A)p(B)+p(A)+p(B)}B\Delta(A, C) \\ & + (-1)^{p(A)+p(B)}A\Delta(B, C) + (-1)^{p(B)}\{A, B, C\}_1,\end{aligned} \tag{2.3}$$

where the last term is the Grassmann-odd Nambu bracket

$$\{A, B, C\}_1 = \epsilon_{\alpha\beta\gamma}\partial_{\theta_\alpha}A\partial_{\theta_\beta}B\partial_{\theta_\gamma}C \tag{2.4}$$

on the Grassmann algebra.

The divergence of the Nambu bracket (2.4) is related with the quantity $\Delta(A, B)$ (2.2) (see also [7])

$$\Delta(A, B) = \frac{1}{2}\partial_{\theta_\alpha}\{\theta_\alpha, A, B\}_1, \tag{2.5}$$

whereas the multiplication on the Grassmann variable θ_α gives the linear odd bracket (1.1)

$$\{A, B\}_1 = \theta_\alpha\{\theta_\alpha, A, B\}_1. \tag{2.6}$$

Note also the following relation between the operator Δ_{-3} (1.4) and odd Nambu bracket (2.4):

$$\Delta_{-3}A = \frac{1}{3!}\partial_{\theta_\alpha}\partial_{\theta_\beta}\{\theta_\alpha, \theta_\beta, A\}_1. \tag{2.7}$$

JHEP00(2002)000

For the operator Δ_{-3} (1.4) there exists the following "Leibniz rule" with respect to the odd Nambu bracket composition:

$$\begin{aligned}\Delta_{-3}(\{A,B,C\}_1) = &- \{\Delta_{-3}A,B,C\}_1 + (-1)^{p(A)}\{A,\Delta_{-3}B,C\}_1\\ &- (-1)^{p(A)+p(B)}\{A,B,\Delta_{-3}C\}_1\\ &+ \epsilon_{\alpha\beta\gamma}\Big[(-1)^{p(A)+1}\Delta\left(\partial_{\theta_\alpha}A,\partial_{\theta_\beta}B\right)\partial_{\theta_\gamma}C\\ &+ (-1)^{p(A)+p(B)}\partial_{\theta_\alpha}A\Delta\left(\partial_{\theta_\beta}B,\partial_{\theta_\gamma}C\right)\\ &+ (-1)^{p(A)p(B)+1}\partial_{\theta_\beta}B\Delta\left(\partial_{\theta_\alpha}A,\partial_{\theta_\gamma}C\right)\Big].\end{aligned} \tag{2.8}$$

It follows from the expression (2.4) the Grassmann parity

$$p(\{A,B,C\}_1) = p(A) + p(B) + p(C) + 1 \pmod 2, \tag{2.9}$$

symmetry properties

$$\{A,B,C\}_1 = -(-1)^{[p(A)+1][p(B)+1]}\{B,A,C\}_1 = -(-1)^{[p(B)+1][p(C)+1]}\{A,C,B\}_1 \tag{2.10}$$

and Jacobi type identity for the odd Nambu bracket

$$\begin{aligned}\{\{A,B,C\}_1,D,E\}_1 &+ (-1)^{[p(A)+1][p(B)+p(C)+p(D)+p(E)]}\{\{B,C,D\}_1,E,A\}_1\\ &+ (-1)^{[p(A)+p(B)][p(C)+p(D)+p(E)+1]}\{\{C,D,E\}_1,A,B\}_1\\ &+ (-1)^{[p(D)+p(E)][p(A)+p(B)+p(C)+1]}\{\{D,E,A\}_1,B,C\}_1\\ &+ (-1)^{[p(E)+1][p(A)+p(B)+p(C)+p(D)]}\{\{E,A,B\}_1,C,D\}_1\\ &+ (-1)^{[p(D)+p(E)][p(A)+p(B)+p(C)+1]+p(B)[p(A)+1]+p(A)}\\ &\times \{\{D,E,B\}_1,A,C\}_1\\ &+ (-1)^{[p(D)+1][p(A)+p(B)+p(C)]+p(D)}\{\{D,A,B\}_1,C,E\}_1\\ &+ (-1)^{[p(A)+1][p(B)+p(C)+p(D)+p(E)]+[p(D)+1]p(E)+p(D)}\\ &\times \{\{B,C,E\}_1,D,A\}_1\\ &+ (-1)^{[p(B)+1][p(C)+p(D)+p(E)]+p(B)}\{\{A,C,D\}_1,E,B\}_1\\ &+ (-1)^{[p(B)+1][p(C)+p(D)+p(E)+1]+[p(D)+1][p(E)+1]}\\ &\times \{\{A,C,E\}_1,D,B\}_1 = 0.\end{aligned} \tag{2.11}$$

Note that the structure of (2.11) is different from the one for the fundamental identity [8].

3. Conclusion

Thus, we constructed the Grassmann-odd Nambu bracket on the Grassmann algebra. The main properties of this bracket are also given.

Acknowledgments

One of the authors (V.A.S.) thanks the administration of the Office of Associate and Federation Schemes of the Abdus Salam ICTP for the kind hospitality at Trieste where this work has been performed.

References

[1] V.A. Soroka, *Degenerate odd Poisson bracket on Grassmann variables*, *Phys. Atom. Nucl.* **63** (2000) 915; `hep-th/9811223`.

[2] V.A. Soroka, *Linear odd Poisson bracket on Grassmann variables*, *Phys. Lett.* **B 451** (1999) 349; `hep-th/9811252`.

[3] V.A. Soroka, *Supersymmetry and the odd Poisson bracket*, *Nucl. Phys.* **101** *(Proc. Suppl.)* (2001) 26; `hep-th/0204018`.

[4] D.V. Soroka and V.A. Soroka, *Lie-Poisson (Kirillov) odd bracket on Grassmann algebra*, Proceedings of the XXIII International Colloquium "Group Theoretical Methods in Physics" (GROUP 23, July 31 – August 5, JINR, Dubna, Russia, 2000) Vol. 1, pp. 106-111, Dubna, 2002.

[5] D.V. Soroka and V.A. Soroka, *Poisson-Lie odd bracket on Grassmann algebra*, *SIGMA* **2** (2006) Paper 036; `hep-th/0603148`.

[6] Y. Nambu, *Generalized Hamiltonian dynamics*, *Phys. Rev.* **D 7** (1973) 2405.

[7] M. Sakakibara, *Notes on the super Nambu bracket*, *Prog. Theor. Phys.* **109** (2003) 305; math-ph/0208040.

[8] L. Takhtajan, *On foundation of the generalized Nambu mechanics*, *Commun. Math. Phys.* **160** (1994) 295.

JHEP00(2002)000